

# DENTALKLINIK Dr.TOKA

H-9400-SOPRON, LACKNERK. STR. 62/B

## SEDATION

STRESS-FREE. RELAXED STATE DURING TREATMENT



This treatment is associated with the least medical complications.  
The most important side effects are the followings:

- feeling sick, vomiting, aspiration (choking on)
- reduced respiratory rate, hypoxia (breathing difficulties)
- drop in blood pressure
- drug allergy
- headache
- violet spot at the puncture site
- even a bigger dose will not help the Patients who refuse the treatment in an extreme way

Please provide exact data for the correct dose:

Name:	<input type="text"/>	Patient's ID:	<input type="text"/>
Body weight:	<input type="text"/> kg	<b>FEE OF THE TREATMENT:</b> First hour of sedation is : <b>320 EUR</b> All inchoate half an hour is : <b>140 EUR</b>	
Height:	<input type="text"/> cm		
Age:	<input type="text"/> years		

Moderate sedation (formerly called "conscious sedation") is carried out by an anaesthesiologist. The anaesthesiologists in charge at our clinic are Dr Árpád Bátorfi and his colleagues. They look forward to answering your questions.

I confirm that I have read and understood the information above. I have informed my doctor about my medical history and I acknowledge the possible risks of the treatment. I hereby consent to the treatment in sedation.

 Sopron

 (day)    (month)    (year)

 Patient

 Doctor

 Nurse

## DENTALKLINIK Dr.TOKA

H-9400-SOPRON, LACKNERK. STR. 62/B

## INFORMATION BOOKLET FOR TREATMENTS CARRIED OUT IN SEDATION

A great result of medicine is that treatments can be carried out in sedation without the patient's feeling the pain. In some cases, some patients are afraid of treatments, but fear, phobia make the accurate and precise work difficult. In this case a treatment in sedation is offered. All over the world the same types of sedation are provided for the unpleasant diagnostic examinations and smaller operations, just like in our dental, implantology and dermatology consulting rooms.

During sedation, moderate sedation we are able to communicate, you can co-operate with us, swallow and other reflexes are still there, but your fear will ease, you will only remember the beginning and the end of the treatment.

Sedation also has an analgesic effect, but at the same time we use local anaesthetic as well. Therefore, in addition to sedation, we also use a local anaesthetic.



## INTRAVENOUS MODERATE SEDATION

for burdensome treatment



## WHAT CAN YOU FEEL DURING THE TREATMENT?

Everybody goes through sedation in different ways

### ADVANTAGES OF INTRAVENOUS MODERATE SEDATION

The advantage is that you can drift away from consciousness and can overcome fear and anxiety. During sedation the anaesthesiologist and his nurses observe the condition of the Patient continually and can adjust the level customized.

Certainly everybody has different experience, there are no two cases alike. It rarely happens that there is no or hardly any sedating effect. Most of the Patients do not remember anything and they are surprised that they are already in the recovery room.


### IMPORTANT TO KNOW:

- 24 hours after sedation it is forbidden to drive, to work at dangerous sites or with dangerous equipment and to drink alcohol.
- You cannot sign any important documents for 24 hours.
- If you arrive at our clinic by car and you go home the same day, you can leave the clinic solely with a company of yours.
- We kindly ask you not to drink or eat anything at least two hours before sedation.
- You will only remember the beginning and the end of the treatment.
- The period of the treatment may be two hours.
- When taking tranquilizers and/or drugs, sedation cannot be applied.
- It is not recommended in the first trimester of pregnancy.
- High blood pressure or heart disease is not a ground for refusal but you should consult with your dentist.
- Because of being monitored after the treatment, you should wait at least for half an hour in our recovery room.

- After receiving the drug, you may feel slightly dizzy and your arms and legs can feel heavy.
- Then everything becomes slowly heavy, you will have apraxia of speech and it can be hardly understood.
- Because of releasing anxiety and inhibitions, some Patients become talkative.
- Some Patients fall asleep but can easily be awakened.
- After the treatment, sense of time declines provisionally.



Please don't hesitate to contact your dentist or our reception for further information.



After the treatment you can relax in the recovery room